

# RT<sup>2</sup> Profiler PCR Array (Rotor-Gene® Format)

## Rabbit Inflammatory Cytokines & Receptors

Cat. no. 330231 PANZ-011ZR

For pathway expression analysis

| Format                                       | For use with the following real-time cyclers |
|----------------------------------------------|----------------------------------------------|
| RT <sup>2</sup> Profiler PCR Array, Format R | Rotor-Gene Q, other Rotor-Gene cyclers       |

### Description

The Rabbit Inflammatory Cytokines & Receptors RT<sup>2</sup> Profiler PCR Array profiles the expression of 84 key genes mediating the inflammatory response. Acute inflammation occurs in response to cell damage due to infection or injury. During this process, cellular and plasma derived factors encourage extravasation, the recruitment of circulating immune cells into the affected tissue. These immune cells in turn up-regulate the expression of inflammatory cytokines that recruit additional immune cells to mount an immune response to any invading organisms and to further promote and eventually resolve the inflammatory response. Chronic inflammation, or expression of these cytokines and receptors at low levels over long periods of time, promotes various pathological conditions including allergies and asthma, cardiovascular system disorders (atherosclerosis), central nervous system disorders (Alzheimer's disease), fibrosis, and rheumatoid arthritis. This array contains genes involved in mediating immune cascade reactions during inflammation. The chemokines, cytokines, and interleukins involved in the inflammatory response are represented as well as their receptors. Profiling the expression of these inflammatory cytokine and receptor genes helps determine the state of and the mechanisms behind inflammation in experimental model systems. Using real-time PCR, research studies can easily and reliably analyze expression of a focused panel of genes related to inflammation with this array.

For further details, consult the *RT<sup>2</sup> Profiler PCR Array Handbook*.

### Shipping and storage

RT<sup>2</sup> Profiler PCR Arrays in the Rotor-Gene format are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products.

---

For long term storage, keep plates at –20°C.

**Note:** Ensure that you have the correct RT<sup>2</sup> Profiler PCR Array format for your real-time cyclers (see table above).

**Note:** Open the package and store the products appropriately immediately on receipt.



---

Sample & Assay Technologies

## Array layout

The 96 real-time assays in the Rotor-Gene format are located in wells 1–96 of the Rotor-Disc™ (plate A1–A12=Rotor-Disc 1–12, plate B1–B12=Rotor-Disc 13–24, etc.). To maintain data analysis compatibility, wells 97–100 do not contain real-time assays but will contain master mix to account for weight balance.

## Gene table: RT<sup>2</sup> Profiler PCR Array

| Position | UniGene  | GenBank      | Symbol       | Description                                                             |
|----------|----------|--------------|--------------|-------------------------------------------------------------------------|
| A01      | Ocu.6220 | XM_002717177 | AIMP1        | Aminoacyl tRNA synthetase complex-interacting multifunctional protein 1 |
| A02      | Ocu.2192 | NM_001082650 | BMP-2        | Bone morphogenetic protein 2                                            |
| A03      | Ocu.2194 | NM_001195723 | BMP4         | Bone morphogenetic protein 4                                            |
| A04      | Ocu.2601 | XM_002724263 | BMP7         | Bone morphogenetic protein 7                                            |
| A05      | Ocu.1921 | NM_001082294 | CCL2         | Chemokine (C-C motif) ligand 2                                          |
| A06      | Ocu.2141 | NM_001082196 | CCL4         | Chemokine (C-C motif) ligand 4                                          |
| A07      | Ocu.7725 | NM_001171124 | CCR1         | Chemokine (C-C motif) receptor 1                                        |
| A08      | Ocu.7516 | NM_001135128 | CCR3         | Chemokine (C-C motif) receptor 3                                        |
| A09      | Ocu.7154 | NM_001171125 | CCR5         | Chemokine (C-C motif) receptor 5 (gene/pseudogene)                      |
| A10      | Ocu.1845 | NM_001082283 | CNTF         | Ciliary neurotrophic factor                                             |
| A11      | Ocu.5753 | NM_001082134 | CX3CR1       | Chemokine (C-X3-C motif) receptor 1                                     |
| A12      | Ocu.7381 | XM_002717002 | CXCL13       | Chemokine (C-X-C motif) ligand 13                                       |
| B01      | Ocu.1917 | NM_001171082 | CXCR1        | Chemokine (C-X-C motif) receptor 1                                      |
| B02      | Ocu.1988 | NM_001171090 | CXCR2        | Chemokine (C-X-C motif) receptor 2                                      |
| B03      | N/A      | XM_002712124 | CXCR4        | Chemokine (C-X-C motif) receptor 4                                      |
| B04      | Ocu.7379 | XM_002722702 | CXCR5        | Chemokine (C-X-C motif) receptor 5                                      |
| B05      | Ocu.7612 | NM_001171257 | GMCSF        | Granulate-macrophage stimulating factor                                 |
| B06      | Ocu.2149 | NM_001081991 | IFNG         | Interferon, gamma                                                       |
| B07      | Ocu.1796 | NM_001082045 | IL10         | Interleukin 10                                                          |
| B08      | Ocu.1795 | NM_001082216 | IL15         | Interleukin 15                                                          |
| B09      | N/A      | XM_002721470 | IL16         | Interleukin 16                                                          |
| B10      | N/A      | XM_002712865 | IL17RA       | Interleukin 17 receptor A                                               |
| B11      | Ocu.7345 | NM_001122940 | IL18         | Interleukin 18 (interferon-gamma-inducing factor)                       |
| B12      | Ocu.7425 | NM_001164863 | IL18R1       | Interleukin 18 receptor 1                                               |
| C01      | Ocu.2080 | NM_001101684 | IL1A         | Interleukin 1, alpha                                                    |
| C02      | Ocu.1993 | NM_001082201 | IL1B         | Interleukin 1, beta                                                     |
| C03      | Ocu.1916 | NM_001082770 | IL1R1        | Interleukin 1 receptor, type I                                          |
| C04      | N/A      | XM_002715977 | IL22RA1      | Interleukin 22 receptor, alpha 1                                        |
| C05      | Ocu.2462 | NM_001163177 | IL4          | Interleukin 4                                                           |
| C06      | N/A      | XM_002711857 | IL4R         | Interleukin 4 receptor                                                  |
| C07      | Ocu.2447 | NM_001082064 | IL6          | Interleukin 6 (interferon, beta 2)                                      |
| C08      | N/A      | XM_002714044 | IL6ST        | Interleukin 6 signal transducer (gp130, oncostatin M receptor)          |
| C09      | N/A      | XM_002714135 | IL7R         | Interleukin 7 receptor                                                  |
| C10      | Ocu.1938 | NM_001082293 | IL8          | Interleukin 8                                                           |
| C11      | Ocu.2072 | NM_001082222 | LOC100009027 | Adiponectin                                                             |
| C12      | Ocu.7378 | XM_002714268 | LOC100144336 | Lymphotoxin beta                                                        |
| D01      | N/A      | XM_002721293 | LOC100338097 | Interferon alpha/beta receptor 2                                        |
| D02      | N/A      | XM_002719359 | LOC100338432 | Chemokine (C-C motif) receptor 7                                        |
| D03      | N/A      | XM_002715435 | LOC100338577 | Colony stimulating factor 1-like                                        |
| D04      | N/A      | XM_002722561 | LOC100338701 | Macrophage migration inhibitory factor                                  |
| D05      | N/A      | XM_002714498 | LOC100339322 | Interleukin 17A-like                                                    |
| D06      | N/A      | XM_002723866 | LOC100339537 | Chemokine (C-C motif) receptor 6-like                                   |
| D07      | N/A      | XM_002714499 | LOC100339570 | Interleukin-17F-like                                                    |
| D08      | N/A      | XM_002710105 | LOC100340820 | Interleukin 9-like                                                      |
| D09      | N/A      | XM_002712966 | LOC100341009 | Tumor necrosis factor ligand superfamily, member 11                     |
|          |          |              | LOC100342    |                                                                         |

| Position | UniGene  | GenBank      | Symbol       | Description                                                     |
|----------|----------|--------------|--------------|-----------------------------------------------------------------|
| D10      | N/A      | XM_002721184 | 491          | Oncostatin M-like                                               |
| D11      | N/A      | XM_002712033 | LOC100343645 | Nicotinamide phosphoribosyltransferase-like                     |
| D12      | N/A      | XM_002717257 | LOC100344172 | Interleukin 21-like                                             |
| E01      | N/A      | XM_002710558 | LOC100345409 | Interleukin 7-like                                              |
| E02      | N/A      | XM_002713106 | LOC100345500 | Chemokine (C-C motif) receptor 8                                |
| E03      | N/A      | XM_002717906 | LOC100345869 | Interleukin 25-like                                             |
| E04      | N/A      | XM_002711541 | LOC100346186 | Chemokine (C-X3-C motif) ligand 1                               |
| E05      | N/A      | XM_002716251 | LOC100347328 | Chemokine (C-C motif) receptor 4                                |
| E06      | N/A      | XM_002720625 | LOC100347606 | Interleukin 5 receptor, alpha-like                              |
| E07      | N/A      | XM_002715338 | LOC100347752 | Interleukin 6 receptor-like                                     |
| E08      | N/A      | XM_002723965 | LOC100347876 | Tumor necrosis factor ligand superfamily, member 14             |
| E09      | N/A      | XM_002707937 | LOC100348362 | Chemokine (C-C motif) ligand 19-like                            |
| E10      | N/A      | XM_002719246 | LOC100348776 | Chemokine (C-C motif) ligand 3-like                             |
| E11      | N/A      | XM_002711248 | LOC100348980 | Interleukin 22                                                  |
| E12      | N/A      | XM_002716291 | LOC100349007 | Interleukin 12A-like                                            |
| F01      | N/A      | XM_002711891 | LOC100349321 | Interleukin 27-like                                             |
| F02      | N/A      | XM_002720089 | LOC100349863 | Chemokine (C-X-C motif) receptor 3                              |
| F03      | N/A      | XM_002715565 | LOC100351924 | Leptin receptor-like                                            |
| F04      | N/A      | XM_002711079 | LOC100352065 | Interleukin 23, alpha subunit p19-like                          |
| F05      | N/A      | XM_002717436 | LOC100353705 | Interleukin 20-like                                             |
| F06      | N/A      | XM_002717112 | LOC100354804 | Chemokine (C-X-C motif) ligand 2-like                           |
| F07      | N/A      | XM_002722190 | LOC100354902 | Interleukin 10 receptor, alpha-like                             |
| F08      | N/A      | XM_002716426 | LOC100355808 | Tumor necrosis factor (ligand) superfamily, member 10-like      |
| F09      | N/A      | XM_002707999 | LOC100356081 | Interleukin 33-like                                             |
| F10      | N/A      | XM_002719135 | LOC100356652 | Colony stimulating factor 3-like                                |
| F11      | N/A      | XM_002710201 | LOC100358075 | Interleukin-5-like                                              |
| F12      | N/A      | XM_002713280 | LOC100358097 | Chemokine (C-C motif) receptor 2                                |
| G01      | N/A      | NM_001256781 | LOC100358388 | CD40 ligand                                                     |
| G02      | N/A      | XM_002710092 | LOC100358676 | Interleukin 13 (predicted)-like                                 |
| G03      | N/A      | NM_001245923 | LOC100358914 | Leukemia inhibitory factor (cholinergic differentiation factor) |
| G04      | Ocu.7553 | NM_001171119 | LTA          | Lymphotoxin alpha (TNF superfamily, member 1)                   |
| G05      | Ocu.2423 | NM_001082194 | SPP1         | Secreted phosphoprotein 1                                       |
| G06      | Ocu.3100 | NM_001082660 | TGFB2        | Transforming growth factor, beta 2                              |
| G07      | Ocu.1971 | NM_001082263 | TNF          | Tumor necrosis factor                                           |
| G08      | Ocu.7986 | XM_002710603 | TNFRSF11B    | Tumor necrosis factor receptor superfamily, member 11b          |
| G09      | Ocu.7332 | NM_001105676 | TNFSF13      | Tumor necrosis factor (ligand) superfamily, member 13           |
| G10      | Ocu.7149 | NM_001099964 | TNFSF13B     | Tumor necrosis factor (ligand) superfamily, member 13b          |
| G11      | Ocu.2137 | NM_001081985 | TNFSF4       | Tumor necrosis factor (ligand) superfamily, member 4            |
| G12      | Ocu.2339 | XM_002714697 | VEGFA        | Vascular endothelial growth factor A                            |
| H01      | Ocu.2090 | NM_001101682 | ACTA2        | Actin, alpha 2, smooth muscle, aorta                            |
| H02      | Ocu.734  | NM_001101683 | ACTB         | Actin, beta                                                     |
| H03      | Ocu.87   | NM_001082253 | GAPDH        | Glyceraldehyde-3-phosphate dehydrogenase                        |

| Position | UniGene  | GenBank      | Symbol           | Description                                     |
|----------|----------|--------------|------------------|-------------------------------------------------|
| H04      | Ocu.1108 | NM_001082277 | LDHA             | Lactate dehydrogenase A                         |
| H05      | N/A      | XM_002723383 | LOC100346<br>936 | Non-POU domain containing, octamer-binding-like |
| H06      | N/A      | SA_00511     | NGDC             | Rabbit Genomic DNA Contamination                |
| H07      | N/A      | SA_00104     | RTC              | Reverse Transcription Control                   |
| H08      | N/A      | SA_00104     | RTC              | Reverse Transcription Control                   |
| H09      | N/A      | SA_00104     | RTC              | Reverse Transcription Control                   |
| H10      | N/A      | SA_00103     | PPC              | Positive PCR Control                            |
| H11      | N/A      | SA_00103     | PPC              | Positive PCR Control                            |
| H12      | N/A      | SA_00103     | PPC              | Positive PCR Control                            |

## Related products

For optimal performance, RT<sup>2</sup> Profiler PCR Arrays should be used together with the RT<sup>2</sup> First Strand Kit for cDNA synthesis and RT<sup>2</sup> SYBR<sup>®</sup> Green qPCR Mastermixes for PCR.

| Product                                                         | Contents                                                                                                 | Cat. no. |
|-----------------------------------------------------------------|----------------------------------------------------------------------------------------------------------|----------|
| RT <sup>2</sup> First Strand Kit (12)                           | Enzymes and reagents for cDNA synthesis                                                                  | 330401   |
| RT <sup>2</sup> SYBR Green ROX <sup>™</sup> FAST Mastermix (2)* | For 2 x 96 assays in 96-well plates; suitable for use with the Rotor-Gene Q and other Rotor-Gene cyclers | 330620   |

\* Larger kit sizes available; please inquire.

RT<sup>2</sup> Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at [www.qiagen.com](http://www.qiagen.com) or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN<sup>®</sup>, Rotor-Gene<sup>®</sup>, Rotor-Disc<sup>™</sup> (QIAGEN Group); ROX<sup>™</sup> (Applied Biosystems or its subsidiaries); SYBR<sup>®</sup> (Molecular Probes, Inc.).

1067688 03/2011 © 2011 QIAGEN, all rights reserved.

[www.qiagen.com](http://www.qiagen.com)

Australia ■ 1-800-243-800

Austria ■ 0800/281010

Belgium ■ 0800-79612

Brazil ■ 0800-557779

Canada ■ 800-572-9613

China ■ 8621-3865-3865

Denmark ■ 80-885945

Finland ■ 0800-914416

France ■ 01-60-920-930

Germany ■ 02103-29-12000

Hong Kong ■ 800 933 965

Ireland ■ 1800 555 049

Italy ■ 800-787980

Japan ■ 03-6890-7300

Korea (South) ■ 080-000-7145

Luxembourg ■ 8002 2076

Mexico ■ 01-800-7742-436

The Netherlands ■ 0800 0229592

Norway ■ 800-18859

Singapore ■ 1800-742-4368

Spain ■ 91-630-7050

Sweden ■ 020-790282

Switzerland ■ 055-254-22-11

UK ■ 01293-422-911

USA ■ 800-426-8157



Sample & Assay Technologies