

RT² Profiler PCR Array (96-Well Format and 384-Well [4 x 96] Format)

Rat Drug Metabolism: Phase II Enzymes

Cat. no. 330231 PARN-069ZA

For pathway expression analysis

| Format | For use with the following real-time cyclers |
|---|--|
| RT ² Profiler PCR Array, Format A | Applied Biosystems® models 5700, 7000, 7300, 7500, 7700, 7900HT, ViiA™ 7 (96-well block); Bio-Rad® models iCycler®, iQ™ 5, MyiQ™, MyiQ2; Bio-Rad/MJ Research Chromo4™; Eppendorf® Mastercycler® ep realplex models 2, 2s, 4, 4s; Stratagene® models Mx3005P®, Mx3000P®; Takara TP-800 |
| RT ² Profiler PCR Array, Format C | Applied Biosystems models 7500 (Fast block), 7900HT (Fast block), StepOnePlus™, ViiA 7 (Fast block) |
| RT ² Profiler PCR Array, Format D | Bio-Rad CFX96™; Bio-Rad/MJ Research models DNA Engine Opticon®, DNA Engine Opticon 2; Stratagene Mx4000® |
| RT ² Profiler PCR Array, Format E | Applied Biosystems models 7900HT (384-well block), ViiA 7 (384-well block); Bio-Rad CFX384™ |
| RT ² Profiler PCR Array, Format F | Roche® LightCycler® 480 (96-well block) |
| RT ² Profiler PCR Array, Format G | Roche LightCycler 480 (384-well block) |
| RT ² Profiler PCR Array, Format H | Fluidigm® BioMark™ |



Sample & Assay Technologies

Description

The Rat Drug Metabolism Phase II Enzymes RT² Profiler PCR Array contains 84 genes involved in the enzymatic processes of drug biotransformation. Phase II drug metabolism enzymes catalyze the conjugation of lipophilic compounds with hydrophilic functional groups or moieties to form water-soluble conjugates that can be cleared from cells and from the body. This array represents genes encoding the seven major classes of phase II drug metabolism enzymes catalyzing such reactions as glutathione conjugation, glucuronidation, sulfation, methylation, amino acid conjugation, epoxidation, and esterification. Using real-time PCR, you can easily and reliably analyze expression of a focused panel of drug metabolism enzymes with this array.

For further details, consult the *RT² Profiler PCR Array Handbook*.


Shipping and storage

RT² Profiler PCR Arrays in formats A, C, D, E, F, and G are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products. RT² Profiler PCR Arrays in format H are shipped on dry ice or blue ice packs.

For long term storage, keep plates at -20°C.

Note: Ensure that you have the correct RT² Profiler PCR Array format for your real-time cycler (see table above).

Note: Open the package and store the products appropriately immediately on receipt.



Array layout (96-well)

For 384-well 4 x 96 PCR arrays, genes are present in a staggered format. Refer to the RT² Profiler PCR Array Handbook for layout.

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
|----------|---------|---------|---------|---------|---------|---------|---------|---------|---------|--------|-------|--------|
| A | A3galt2 | A4galt | Aanat | Acsl1 | Acsl3 | Acsl4 | Acsn3 | Agxt | Alg5 | As3mt | Asmt | Bcat |
| B | Ccbl1 | Ces1d | Ces1e | Ces2c | Ces5a | Chst7 | Comt | Ddost | Eef1b2 | Ephx1 | Ephx2 | Galnt1 |
| C | Galnt3 | Galnt4 | Gamt | Gcnt1 | Gcnf2 | Glyat | Gnmt | Gsta3 | Gsta4 | Gstk1 | Gstm1 | Gstm2 |
| D | Gstm3 | Gstm4 | Gstm5 | Gsto1 | Gsto2 | Gstp1 | Gstt1 | Gstt2 | Has1 | Hnmt | Inmt | Mgal1 |
| E | Mgal2 | Mgst1 | Mgst2 | Mgst3 | Nat1 | Nat2 | Nnmt | Nqo1 | Nqo2 | Pomgn1 | Ptges | Ptges2 |
| F | Sat1 | Sult1a1 | Sult1b1 | Sult1c2 | Sult1e1 | Sult2b1 | Sult4a1 | Sult5a1 | Sult6b1 | Tst | Ugcg | |
| G | Ugt1a1 | Ugt1a5 | Ugt1a6 | Ugt1a9 | Ugt2a1 | Ugt2a3 | Ugt2b | Ugt2b1 | Ugt2b5 | Ugt8 | Xdh | |
| H | Actb | B2m | Hprt1 | Ldha | Rplp1 | RGDC | RTC | RTC | PPC | PPC | PPC | PPC |

Gene table: RT² Profiler PCR Array

| Position | UniGene | GenBank | Symbol | Description |
|----------|-----------|--------------|---------|--|
| A01 | Rn.37737 | NM_138524 | A3galt2 | Alpha 1,3-galactosyltransferase 2 |
| A02 | Rn.48584 | NM_022240 | A4galt | Alpha 1,4-galactosyltransferase |
| A03 | Rn.88180 | NM_012818 | Aanat | Arylalkylamine N-acetyltransferase |
| A04 | Rn.6215 | NM_012820 | Acsl1 | Acyl-CoA synthetase long-chain family member 1 |
| A05 | Rn.54820 | NM_057107 | Acsl3 | Acyl-CoA synthetase long-chain family member 3 |
| A06 | Rn.87821 | NM_053623 | Acsl4 | Acyl-CoA synthetase long-chain family member 4 |
| A07 | Rn.88644 | NM_033231 | Acsn3 | Acyl-CoA synthetase medium-chain family member 3 |
| A08 | Rn.9931 | NM_030656 | Agxt | Alanine-glyoxylate aminotransferase |
| A09 | Rn.155539 | NM_001025407 | Alg5 | Asparagine-linked glycosylation 5, dolichyl-phosphate beta-glucosyltransferase homolog (S. cerevisiae) |
| A10 | Rn.95453 | NM_080890 | As3mt | Arsenic (+3 oxidation state) methyltransferase |
| A11 | Rn.10768 | NM_144759 | Asmt | Acetylserotonin O-methyltransferase |
| A12 | Rn.11129 | NM_017300 | Bcat | Bile acid Coenzyme A: amino acid N-acyltransferase (glycine N-choloyltransferase) |
| B01 | Rn.110564 | NM_001013164 | Ccbl1 | Cysteine conjugate-beta lyase, cytoplasmic |
| B02 | Rn.201285 | NM_133295 | Ces1d | Carboxylesterase 1D |
| B03 | Rn.161717 | NM_031565 | Ces1e | Carboxylesterase 1E |
| B04 | Rn.14535 | NM_133586 | Ces2c | Carboxylesterase 2C |
| B05 | Rn.61848 | NM_001012056 | Ces5a | Carboxylesterase 5A |
| B06 | Rn.153551 | NM_207600 | Chst7 | Carbohydrate (N-acetylglucosamine 6-O) sulfotransferase 7 |
| B07 | Rn.220 | NM_012531 | Comt | Catechol-O-methyltransferase |
| B08 | Rn.102842 | NM_001012104 | Ddost | Dolichyl-diphosphooligosaccharide-protein glycosyltransferase |
| B09 | Rn.3910 | NM_001108799 | Eef1b2 | Eukaryotic translation elongation factor 1 beta 2 |
| B10 | Rn.3603 | NM_012844 | Ephx1 | Epoxide hydrolase 1, microsomal |
| B11 | Rn.54495 | NM_022936 | Ephx2 | Epoxide hydrolase 2, cytoplasmic |
| B12 | Rn.10266 | NM_024373 | Galnt1 | UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase 1 (GalNAc-T1) |
| C01 | Rn.78014 | NM_001015032 | Galnt3 | UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase 3 (GalNAc-T3) |
| C02 | Rn.207152 | NM_001025053 | Galnt4 | UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase 4 (GalNAc-T4) |
| C03 | Rn.33890 | NM_012793 | Gamt | Guanidinoacetate N-methyltransferase |
| C04 | Rn.162516 | NM_022276 | Gcnt1 | Glucosaminyl (N-acetyl) transferase 1, core 2 (beta-1, 6-N-acetylglucosaminyltransferase) |
| C05 | Rn.8807 | NM_001001511 | Gcnt2 | Glucosaminyl (N-acetyl) transferase 2, I-branching enzyme |
| C06 | Rn.21872 | NM_001009648 | Glyat | Glycine-N-acyltransferase |
| C07 | Rn.11142 | NM_017084 | Gnmt | Glycine N-methyltransferase |
| C08 | Rn.10460 | NM_031509 | Gsta3 | Glutathione S-transferase A3 |
| C09 | Rn.57528 | NM_001106840 | Gsta4 | Glutathione S-transferase alpha 4 |
| C10 | Rn.109452 | NM_181371 | Gstk1 | Glutathione S-transferase kappa 1 |
| C11 | Rn.202944 | NM_017014 | Gstm1 | Glutathione S-transferase mu 1 |
| C12 | Rn.625 | NM_177426 | Gstm2 | Glutathione S-transferase mu 2 |
| D01 | Rn.6036 | NM_020540 | Gstm3 | Glutathione S-transferase mu 3 |
| D02 | Rn.209043 | NM_001024304 | Gstm4 | Glutathione S-transferase mu 4 |
| D03 | Rn.9158 | NM_172038 | Gstm5 | Glutathione S-transferase, mu 5 |

| Position | UniGene | GenBank | Symbol | Description |
|----------|-----------|--------------|---------|--|
| D04 | Rn.25166 | NM_001007602 | Gsto1 | Glutathione S-transferase omega 1 |
| D05 | Rn.206114 | NM_001012071 | Gsto2 | Glutathione S-transferase omega 2 |
| D06 | Rn.87063 | NM_012577 | Gstp1 | Glutathione S-transferase pi 1 |
| D07 | Rn.11122 | NM_053293 | Gstt1 | Glutathione S-transferase theta 1 |
| D08 | Rn.87212 | NM_012796 | Gst2 | Glutathione S-transferase, theta 2 |
| D09 | Rn.91138 | NM_172323 | Has1 | Hyaluronan synthase 1 |
| D10 | Rn.13145 | NM_031044 | Hnm1 | Histamine N-methyltransferase |
| D11 | Rn.19133 | NM_001109022 | Inmt | Indolethylamine N-methyltransferase |
| D12 | Rn.2712 | NM_030861 | Mgat1 | Mannosyl (alpha-1,3)-glycoprotein beta-1,2-N-acetylglucosaminyltransferase |
| E01 | Rn.2342 | NM_053604 | Mgat2 | Mannosyl (alpha-1,6)-glycoprotein beta-1,2-N-acetylglucosaminyltransferase |
| E02 | Rn.2580 | NM_134349 | Mgst1 | Microsomal glutathione S-transferase 1 |
| E03 | Rn.7854 | NM_001106430 | Mgst2 | Microsomal glutathione S-transferase 2 |
| E04 | Rn.1916 | XM_213943 | Mgst3 | Microsomal glutathione S-transferase 3 |
| E05 | Rn.37420 | NM_001037315 | Nat1 | N-acetyltransferase 1 |
| E06 | Rn.198370 | NM_053854 | Nat2 | N-acetyltransferase 2 |
| E07 | Rn.15755 | NM_001106819 | Nnmt | Nicotinamide N-methyltransferase |
| E08 | Rn.11234 | NM_017000 | Nqo1 | NAD(P)H dehydrogenase, quinone 1 |
| E09 | Rn.7882 | NM_001004214 | Nqo2 | NAD(P)H dehydrogenase, quinone 2 |
| E10 | Rn.162088 | NM_001007747 | Pomgn1 | Protein O-linked mannose beta 1,2-N-acetylglucosaminyltransferase |
| E11 | Rn.7730 | NM_021583 | Ptg5 | Prostaglandin E synthase |
| E12 | Rn.39290 | NM_001107832 | Ptg5e2 | Prostaglandin E synthase 2 |
| F01 | Rn.107986 | NM_001007667 | Sat1 | Spermidine/spermine N1-acetyl transferase 1 |
| F02 | Rn.1507 | NM_031834 | Sult1a1 | Sulfotransferase family, cytosolic, 1A, phenol-preferring, member 1 |
| F03 | Rn.1183 | NM_022513 | Sult1b1 | Sulfotransferase family, cytosolic, 1B, member 1 |
| F04 | Rn.220698 | NM_133547 | Sult1c2 | Sulfotransferase family, cytosolic, 1C, member 2 |
| F05 | Rn.9937 | NM_031732 | Sult1c3 | Sulfotransferase family, cytosolic, 1C, member 3 |
| F06 | Rn.32104 | NM_012883 | Sult1e1 | Sulfotransferase family 1E, estrogen-preferring, member 1 |
| F07 | Rn.100005 | NM_001039665 | Sult2b1 | Sulfotransferase family, cytosolic, 2B, member 1 |
| F08 | Rn.162965 | NM_031641 | Sulta4l | Sulfotransferase family 4A, member 1 |
| F09 | Rn.107747 | NM_001106194 | Sult5a1 | Sulfotransferase family 5A, member 1 |
| F10 | Rn.155063 | XM_578623 | Sult6b1 | Sulfotransferase family, cytosolic, 6B, member 1 |
| F11 | Rn.6360 | NM_012808 | Tst | Thiosulfate sulfurtransferase |
| F12 | Rn.24091 | NM_031795 | Ugcg | UDP-glucose ceramide glucosyltransferase |
| G01 | Rn.26489 | NM_012683 | Ugt1a1 | UDP glucuronosyltransferase 1 family, polypeptide A1 |
| G02 | Rn.26489 | NM_001039549 | Ugt1a5 | UDP glucuronosyltransferase 1 family, polypeptide A5 |
| G03 | Rn.26489 | NM_057105 | Ugt1a6 | UDP glucuronosyltransferase 1 family, polypeptide A6 |
| G04 | Rn.26489 | NM_175846 | Ugt1a8 | UDP glycosyltransferase 1 family, polypeptide A8 |
| G05 | Rn.26489 | NM_201425 | Ugt1a9 | UDP glucuronosyltransferase 1 family, polypeptide A9 |
| G06 | Rn.138121 | NM_022228 | Ugt2a1 | UDP glucuronosyltransferase 2 family, polypeptide A1 |
| G07 | Rn.124736 | NM_001135869 | Ugt2a3 | UDP glucuronosyltransferase 2 family, polypeptide A3 |
| G08 | Rn.2521 | NM_031533 | Ugt2b2 | UDP glycosyltransferase 2 family, polypeptide B |
| G09 | Rn.9969 | NM_173295 | Ugt2b1 | UDP glucuronosyltransferase 2 family, polypeptide B1 |
| G10 | Rn.217178 | NM_153314 | Ugt2b5 | UDP glucuronosyltransferase 2 family, polypeptide B5 |
| G11 | Rn.9744 | NM_019276 | Ugt8 | UDP glycosyltransferase 8 |
| G12 | Rn.202951 | NM_017154 | Xdh | Xanthine dehydrogenase |
| H01 | Rn.94978 | NM_031144 | Actb | Actin, beta |
| H02 | Rn.1868 | NM_012512 | B2m | Beta-2 microglobulin |
| H03 | Rn.47 | NM_012583 | Hprt1 | Hypoxanthine phosphoribosyltransferase 1 |
| H04 | Rn.107896 | NM_017025 | Ldha | Lactate dehydrogenase A |
| H05 | Rn.973 | NM_001007604 | Rplp1 | Ribosomal protein, large, P1 |
| H06 | N/A | U26919 | RGDC | Rat Genomic DNA Contamination |
| H07 | N/A | SA_00104 | RTC | Reverse Transcription Control |
| H08 | N/A | SA_00104 | RTC | Reverse Transcription Control |
| H09 | N/A | SA_00104 | RTC | Reverse Transcription Control |
| H10 | N/A | SA_00103 | PPC | Positive PCR Control |
| H11 | N/A | SA_00103 | PPC | Positive PCR Control |
| H12 | N/A | SA_00103 | PPC | Positive PCR Control |

Related products

For optimal performance, RT² Profiler PCR Arrays should be used together with the RT² First Strand Kit for cDNA synthesis and RT2 SYBR® Green qPCR Mastermixes for PCR.

| Product | Contents | Cat. no. |
|--|---|----------|
| RT ² First Strand Kit (12) | Enzymes and reagents for cDNA synthesis | 330401 |
| RT ² SYBR Green qPCR Mastermix (2)* | For 2 x 96 assays in 96-well plates; suitable for use with real-time cyclers that do not require a reference dye, including: Bio-Rad models CFX96, CFX384, DNA Engine Opticon 2; Bio-Rad/MJ Research Chromo4; Roche LightCycler 480 (96-well and 384-well); all other cyclers | 330500 |
| RT ² SYBR Green ROX™ qPCR Mastermix (2)* | For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Applied Biosystems models 5700, 7000, 7300, 7500 [Standard and FAST], 7700, 7900HT 96-well block [Standard and FAST] and 384-well block, StepOnePlus; Eppendorf Mastercycler ep realplex models 2, 2S, 4, 4S; Stratagene models Mx3000P, Mx3005P, Mx4000; Takara TP-800 | 330520 |
| RT ² SYBR Green Fluor qPCR Mastermix (2)* | For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Bio-Rad models iCycler, iQ5, MyiQ, MyiQ2 | 330510 |

* Larger kit sizes available; please inquire.

RT² Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen.com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN® (QIAGEN Group); Applied Biosystems®, ViiA™, StepOnePlus™, ROX™ (Applied Biosystems or its subsidiaries); Bio-Rad®, iCycler®, iQ™, MyIQ™, Chromo4™, CFX96™, DNA Engine Opticon®, CFX384™ (Bio-Rad Laboratories, Inc.) Stratagene®, Mx3005P®, Mx3000P®, Mx4000® (Stratagene); Eppendorf®, Mastercycler® (Eppendorf AG); Roche®, LightCycler® (Roche Group); Fluidigm® BioMark™ (Fluidigm Corporation); SYBR® (Molecular Probes, Inc.).

1066029 03/2011 © 2011 QIAGEN, all rights reserved.

www.qiagen.com

Canada ■ 800-572-9613

China ■ 8621-3865-3865

Denmark ■ 80-885945

Australia ■ 1-800-243-800

Austria ■ 0800/281010

Belgium ■ 0800-79612

Brazil ■ 0800-557779

Ireland ■ 1800 555 049

Italy ■ 800-787980

Japan ■ 03-6890-7300

Finland ■ 0800-914416

France ■ 01-60-920-930

Germany ■ 02103-29-12000

Hong Kong ■ 800 933 965

Norway ■ 800-18859

Singapore ■ 1800-742-4368

Spain ■ 91-630-7050

Sweden ■ 020-790282

Luxembourg ■ 8002 2076

Korea (South) ■ 080-000-7145

Mexico ■ 01-800-7742-436

The Netherlands ■ 0800 0229592

UK ■ 01293-422-911

USA ■ 800-426-8157



Sample & Assay Technologies